

Universal Bell Correlations Do Not Exist

Cole A. Graham¹ **William M. Hoza**²

December 4, 2016

CS395T – Quantum Complexity Theory

¹Department of Mathematics, Stanford University

²Department of Computer Science, UT Austin

Quantum nonlocality

- ▶ Recall **Bell's theorem**: Entanglement allows interactions that can't be simulated using shared randomness / hidden variables

Quantum nonlocality

- ▶ Recall **Bell's theorem**: Entanglement allows interactions that can't be simulated using shared randomness / hidden variables
- ▶ Recall the **no-communication theorem**: Entanglement can't be used to send signals

Quantum nonlocality


- ▶ Recall **Bell's theorem**: Entanglement allows interactions that can't be simulated using shared randomness / hidden variables
- ▶ Recall the **no-communication theorem**: Entanglement can't be used to send signals
- ▶ **Contradictory?**

Alice




Bob


PR box



PR box




PR box



$$(a, b) = \begin{cases} (0, xy) & \text{with probability } 1/2 \\ (1, 1 - xy) & \text{with probability } 1/2 \end{cases}$$


PR box



$$(a, b) = \begin{cases} (0, xy) & \text{with probability } 1/2 \\ (1, 1 - xy) & \text{with probability } 1/2 \end{cases}$$

- ▶ Cannot be used to **communicate**


PR box




$$(a, b) = \begin{cases} (0, xy) & \text{with probability } 1/2 \\ (1, 1 - xy) & \text{with probability } 1/2 \end{cases}$$

- ▶ Cannot be used to **communicate**
- ▶ But can be used to **win CHSH game**: $a + b = xy \pmod{2}$

Correlation box




Correlation box



- ▶ A *correlation box* is a map

$$\text{Cor} : X \times Y \rightarrow \{\mu : \mu \text{ is a probability distribution over } A \times B\}$$

Correlation box




- ▶ A *correlation box* is a map

$$\text{Cor} : X \times Y \rightarrow \{\mu : \mu \text{ is a probability distribution over } A \times B\}$$

- ▶ Assume X, Y, A, B are countable

Correlation box




- ▶ A *correlation box* is a map

$$\text{Cor} : X \times Y \rightarrow \{\mu : \mu \text{ is a probability distribution over } A \times B\}$$

- ▶ Assume X, Y, A, B are countable
- ▶ Abuse notation and write $\text{Cor} : X \times Y \rightarrow A \times B$

Distributed sampling problems



- ▶ Can think of a correlation box as a *distributed sampling problem* – the **problem of simulating the box**

Distributed sampling complexity classes

- ▶ **SR**: class of correlation boxes that can be simulated using **just shared randomness**

Distributed sampling complexity classes

- ▶ **SR**: class of correlation boxes that can be simulated using **just shared randomness**
- ▶ **Q**: class of correlation boxes that can be simulated using **shared randomness + arbitrary bipartite quantum state**

Distributed sampling complexity classes

- ▶ **SR**: class of correlation boxes that can be simulated using **just shared randomness**
- ▶ **Q**: class of correlation boxes that can be simulated using **shared randomness + arbitrary bipartite quantum state**
- ▶ Obviously **SR** \subseteq **Q**

Distributed sampling complexity classes

- ▶ **SR**: class of correlation boxes that can be simulated using **just shared randomness**
- ▶ **Q**: class of correlation boxes that can be simulated using **shared randomness + arbitrary bipartite quantum state**
- ▶ Obviously **SR** \subseteq **Q**
- ▶ Bell's theorem: **SR** \neq **Q**

Distributed sampling complexity classes (2)

- ▶ **NS**: class of non-signalling correlation boxes

Distributed sampling complexity classes (2)

- ▶ **NS**: class of non-signalling correlation boxes
- ▶ No-communication theorem: $\mathbf{Q} \subseteq \mathbf{NS}$

Distributed sampling complexity classes (2)

- ▶ **NS**: class of non-signalling correlation boxes
- ▶ No-communication theorem: $\mathbf{Q} \subseteq \mathbf{NS}$
- ▶ Tsierelson bound: $\text{PR} \notin \mathbf{Q}$, so $\mathbf{Q} \neq \mathbf{NS}$

Bell pair

- ▶ Goal: Understand **Q**

Bell pair

- ▶ Goal: Understand **Q**
- ▶ Baby step: Understand **BELL**: class of correlation boxes that can be simulated using **shared randomness** + $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ + **projective measurements**

Bell pair

- ▶ Goal: Understand **Q**
- ▶ Baby step: Understand **BELL**: class of correlation boxes that can be simulated using **shared randomness** + $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ + **projective measurements**
- ▶ **SR** \subsetneq **BELL** \subsetneq **Q**

Toner-Bacon theorem

- ▶ Theorem (Toner, Bacon '03): **BELL** can be simulated using shared randomness + 1 bit of communication

Toner-Bacon theorem

- ▶ Theorem (Toner, Bacon '03): **BELL** can be simulated using **shared randomness + 1 bit of communication**
- ▶ This is an **upper bound** on the power of **BELL**

Toner-Bacon theorem

- ▶ Theorem (Toner, Bacon '03): **BELL** can be simulated using **shared randomness + 1 bit of communication**
- ▶ This is an **upper bound** on the power of **BELL**
- ▶ **Loose** upper bound, since **BELL** \subseteq **NS**


PR box is **BELL**-hard

- ▶ Theorem (Cerf et al. '05): **BELL** can be simulated using shared randomness + 1 PR box

PR box is **BELL**-hard

- ▶ Theorem (Cerf et al. '05): **BELL** can be simulated using **shared randomness + 1 PR box**
- ▶ In other words, PR is **BELL**-hard with respect to **1-query reductions**

Distributed sampling complexity zoo



- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box

- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box



- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box



- ▶ Theorem:

- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box



- ▶ Theorem:

- ▶ Suppose $\text{Cor} : X \times Y \rightarrow A \times B$ is in **Q**; X, Y countable; A, B finite

- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box



- ▶ Theorem:

- ▶ Suppose $\text{Cor} : X \times Y \rightarrow A \times B$ is in **Q**; X, Y countable; A, B finite
- ▶ Then there exists a **binary** correlation box in **BELL** that does not reduce to Cor

- ▶ Theorem: There does not exist a finite-alphabet **BELL**-complete correlation box




- ▶ Theorem:


- ▶ Suppose $\text{Cor} : X \times Y \rightarrow A \times B$ is in **Q**; X, Y countable; A, B finite
- ▶ Then there exists a **binary** correlation box in **BELL** that does not reduce to Cor



Distributed sampling complexity zoo (2)




Biased CHSH game




- ▶ Goal: $a + b = xy \pmod{2}$

Biased CHSH game




- ▶ Goal: $a + b = xy \pmod{2}$
- ▶ Inputs x, y are chosen independently at random

Biased CHSH game



- ▶ Goal: $a + b = xy \pmod{2}$
- ▶ Inputs x, y are chosen independently at random
- ▶ y is uniform, x is biased: $\Pr[x = 1] = p \in [1/2, 1]$


Biased CHSH game



- ▶ Goal: $a + b = xy \pmod{2}$
- ▶ Inputs x, y are chosen independently at random
- ▶ y is uniform, x is biased: $\Pr[x = 1] = p \in [1/2, 1]$
- ▶ Theorem (Lawson, Linden, Popescu '10): Optimal quantum strategy can be implemented in **BELL**, wins with probability

$$f(p) \stackrel{\text{def}}{=} \frac{1}{2} + \frac{1}{2} \sqrt{p^2 + (1-p)^2}$$

Quantum value of biased CHSH game



Affine functions from reductions

- ▶ Let $S_p \in \mathbf{BELL}$ be optimal quantum strategy

Affine functions from reductions

- ▶ Let $S_p \in \mathbf{BELL}$ be optimal quantum strategy
- ▶ Assume there is a reduction from S_p to Cor

Affine functions from reductions

- ▶ Let $S_p \in \mathbf{BELL}$ be optimal quantum strategy
- ▶ Assume there is a reduction from S_p to Cor
- ▶ Probability that reduction wins biased CHSH game is of form

$$\frac{1-p}{2}P_{00} + \frac{p}{2}P_{10} + \frac{1-p}{2}P_{01} + \frac{p}{2}P_{11}$$

Affine functions from reductions

- ▶ Let $S_p \in \mathbf{BELL}$ be optimal quantum strategy
- ▶ Assume there is a reduction from S_p to Cor
- ▶ Probability that reduction wins biased CHSH game is of form

$$\frac{1-p}{2}P_{00} + \frac{p}{2}P_{10} + \frac{1-p}{2}P_{01} + \frac{p}{2}P_{11}$$

- ▶ Affine function of p , for fixed reduction

Countably many affine functions

- ▶ **Fix shared randomness** without decreasing win probability

Countably many affine functions

- ▶ **Fix shared randomness** without decreasing win probability
- ▶ Recall $\text{Cor} \in \mathbf{Q}$

Countably many affine functions

- ▶ Fix shared randomness without decreasing win probability
- ▶ Recall $\text{Cor} \in \mathbf{Q}$
- ▶ Win probability still exactly $f(p)$ (\mathbf{Q} is closed under reductions)

Countably many affine functions

- ▶ **Fix shared randomness** without decreasing win probability
- ▶ Recall $\text{Cor} \in \mathbf{Q}$
- ▶ Win probability still exactly $f(p)$ (\mathbf{Q} is closed under reductions)
- ▶ Recall $\text{Cor} : X \times Y \rightarrow A \times B$ with X, Y countable, A, B finite

Countably many affine functions

- ▶ **Fix shared randomness** without decreasing win probability
- ▶ Recall $\text{Cor} \in \mathbf{Q}$
- ▶ Win probability still exactly $f(p)$ (\mathbf{Q} is closed under reductions)
- ▶ Recall $\text{Cor} : X \times Y \rightarrow A \times B$ with X, Y countable, A, B finite
- ▶ Only **countably many** deterministic reductions!

Countably many affine functions

- ▶ Fix shared randomness without decreasing win probability
- ▶ Recall $\text{Cor} \in \mathbf{Q}$
- ▶ Win probability still exactly $f(p)$ (\mathbf{Q} is closed under reductions)
- ▶ Recall $\text{Cor} : X \times Y \rightarrow A \times B$ with X, Y countable, A, B finite
- ▶ Only countably many deterministic reductions!
- ▶ Countably many affine functions, so $\exists p$ where all the affine functions disagree with $f(p)$

Approximate simulations

- ▶ Theorem:

Approximate simulations

- ▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in \mathbf{Q} ; X, Y, A, B finite

Approximate simulations

- ▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in$ **BELL** such that

Approximate simulations

▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in \mathbf{BELL}$ such that
- ▶ If there is a k -query ϵ -error reduction from Cor_1 to Cor_2 , then

$$k^4 \cdot (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k} \geq \Omega(1/\epsilon)$$

Approximate simulations

▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in \mathbf{BELL}$ such that
- ▶ If there is a k -query ϵ -error reduction from Cor_1 to Cor_2 , then

$$k^4 \cdot (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k} \geq \Omega(1/\epsilon)$$

- ▶ Upper bound: $\forall \epsilon > 0, \exists \text{Cor}_2 : [T] \times [T] \rightarrow \{0, 1\} \times \{0, 1\}$ such that

Approximate simulations

▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in \mathbf{BELL}$ such that
- ▶ If there is a k -query ϵ -error reduction from Cor_1 to Cor_2 , then

$$k^4 \cdot (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k} \geq \Omega(1/\epsilon)$$

- ▶ Upper bound: $\forall \epsilon > 0, \exists \text{Cor}_2 : [T] \times [T] \rightarrow \{0, 1\} \times \{0, 1\}$ such that
 - ▶ $T \leq O(1/\epsilon^4)$

Approximate simulations

▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in \mathbf{BELL}$ such that
- ▶ If there is a k -query ϵ -error reduction from Cor_1 to Cor_2 , then

$$k^4 \cdot (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k} \geq \Omega(1/\epsilon)$$

- ▶ Upper bound: $\forall \epsilon > 0, \exists \text{Cor}_2 : [T] \times [T] \rightarrow \{0, 1\} \times \{0, 1\}$ such that
 - ▶ $T \leq O(1/\epsilon^4)$
 - ▶ $\text{Cor}_2 \in \mathbf{BELL}$

Approximate simulations

▶ Theorem:

- ▶ Suppose $\text{Cor}_2 : X \times Y \rightarrow A \times B$ is in **Q**; X, Y, A, B finite
- ▶ Then \exists binary correlation box $\text{Cor}_1 \in \mathbf{BELL}$ such that
- ▶ If there is a k -query ε -error reduction from Cor_1 to Cor_2 , then

$$k^4 \cdot (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k} \geq \Omega(1/\varepsilon)$$

▶ Upper bound: $\forall \varepsilon > 0, \exists \text{Cor}_2 : [T] \times [T] \rightarrow \{0, 1\} \times \{0, 1\}$
such that

- ▶ $T \leq O(1/\varepsilon^4)$
- ▶ $\text{Cor}_2 \in \mathbf{BELL}$
- ▶ For every $\text{Cor}_1 \in \mathbf{BELL}$, there is a 1-query ε -error reduction from Cor_1 to Cor_2

Conclusions

- ▶ Is there a **countable**-alphabet **BELL**-complete correlation box?

Conclusions

- ▶ Is there a **countable**-alphabet **BELL**-complete correlation box?
- ▶ What is the right relationship between $|X|, |Y|, |A|, |B|, k, \varepsilon$?

Conclusions

- ▶ Is there a **countable**-alphabet **BELL**-complete correlation box?
- ▶ What is the right relationship between $|X|, |Y|, |A|, |B|, k, \epsilon$?

▶ Thanks for listening!
Questions?

- ▶ This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE-1610403.
- ▶ Cole Graham gratefully acknowledges the support of the Fannie and John Hertz Foundation.